

Name _____ DOB ____/____/____

Severity Classification ☐ Intermittent ☐ Mild Persistent ☐ Moderate Persistent ☐ Severe Persistent

Asthma Triggers (list) _____

Peak Flow Meter Personal Best _____

Green Zone: Doing Well

Symptoms: Breathing is good – No cough or wheeze – Can work and play – Sleeps well at night

Peak Flow Meter _____ (more than 80% of personal best)

Control Medicine(s)	Medicine	How much to take	When and how often to take it
	_____	_____	_____
	_____	_____	_____

Physical Activity ☐ Use albuterol/levalbuterol ____ puffs, 15 minutes before activity
☐ with all activity ☐ when you feel you need it

Yellow Zone: Caution

Symptoms: Some problems breathing – Cough, wheeze, or chest tight – Problems working or playing – Wake at night

Peak Flow Meter _____ to _____ (between 50% and 79% of personal best)

Quick-relief Medicine(s) ☐ Albuterol/levalbuterol ____ puffs, every 4 hours as needed

Control Medicine(s) ☐ Continue Green Zone medicines
☐ Add _____ ☐ Change to _____

You should feel better within 20–60 minutes of the quick-relief treatment. If you are getting worse or are in the Yellow Zone for more than 24 hours, THEN follow the instructions in the RED ZONE and call the doctor right away!

Red Zone: Get Help Now!

Symptoms: Lots of problems breathing – Cannot work or play – Getting worse instead of better – Medicine is not helping

Peak Flow Meter _____ (less than 50% of personal best)

Take Quick-relief Medicine NOW! ☐ Albuterol/levalbuterol ____ puffs, _____ (how frequently)

Call 911 immediately if the following danger signs are present

- Trouble walking/talking due to shortness of breath
- Lips or fingernails are blue
- Still in the red zone after 15 minutes

Emergency Contact Name _____ Phone (____) _____ - _____

Healthcare Provider Name _____ Phone (____) _____ - _____